

## What You'll Learn

- Expand your understanding of solving equations.
- Model and solve problems using linear equations.
- Investigate the properties of inequalities.
- Explain and illustrate strategies to solve linear inequalities.

# Why It's Important

Linear equations and inequalities are used by

- nurses, home health aides, and medical assistants, to take temperatures and blood pressures, and set up equipment
- purchasing agents and buyers, to find the best merchandise at the lowest price for their employers, and stay aware of changes in the marketplace


## **Key Words**

inverse operations variable inequality

# 6.1 Skill Builder

# **Order of Operations**

We use this order of operations to evaluate expressions with more than one operation.




## Check

1. In each expression, circle what you will do first.

<b>a)</b> -7 + 2 × (-3)	Add	Multiply		
<b>b)</b> 3 × (-10 ÷ 2) - (-4)	Multiply	Divide	Subtract	
<b>c)</b> $19 - 4 \times 3^2 \div 6$	Subtract	Multiply	Power	Divide
<b>d)</b> −30 ÷ 5 − 10 × 2	Divide	Subtract	Multiply	


2. Evaluate.



The Distributive PropertyTo multiply  $5 \times (3 + 4)$ , we can:• Add 3 + 4 in the brackets, then multiply the sum by 5: $5 \times (3 + 4)$  $= 5 \times 7$ = 35OR• Multiply each number in the brackets by 5, then add: $5 \times (3 + 4) = 5 \times 3 + 5 \times 4$ = 15 + 20= 35We can use the distributive property to write this expression as a sum of terms:7(a + b) = 7a + 7b


### Check

1. Expand.



# 6.1 Solving Equations by Using Inverse Operations

# **FOCUS** Model a problem with a linear equation, and solve the equation pictorially and symbolically.



**Inverse operations** undo each other's results. For example: addition and subtraction are inverse operations.

# **Example 1** Writing Then Solving One-Step Equations


Write then solve an equation to find each number. Verify the solution.

```
a) A number plus 5 is 20. b) Four times a number is -32.
```

### Solution

a) Let x represent the number. Then, x plus 5 is 20. The equation is: x + 5 = 20To solve the equation, apply the inverse operations. x + 5 = 20Undo the addition. Subtract 5 from each side. x + 5 - 5 = 20 - 5x = 15To verify the solution, substitute x = 15 into x + 5 = 20. 15 + 5 = 20, so the solution is correct. **b)** Let *n* represent the number. Then, 4 times *n* is -32. The equation is: 4n = -32To solve the equation, apply the inverse operations. 4n = -32Undo the multiplication. Divide each side by 4.  $\frac{4n}{2} = \frac{-32}{2}$ n = -8To verify the solution, substitute n = -8 into 4n = -32. 4(-8) = -32, so the solution is correct.

**1.** Let *n* represent a number. Two less than a number is 10. What is the number?



# **Example 2** Solving a Two-Step Equation

Solve, then verify each equation.

**a)** 3x + 4 = -5 **b)** 2(-2 + w) = 18

### Solution

a) Perform the inverse operations in reverse order. 3x + 4 = -5Subtract 4 from each side. 3x + 4 - 4 = -5 - 4 3x = -9Divide each side by 3.  $\frac{3x}{3} = \frac{-9}{3}$  x = -3

To verify the solution, substitute x = -3 into 3x + 4 = -5. Left side = 3x + 4Right side = -5= 3(-3) + 4= -9 + 4

Since the left side equals the right side, x = -3 is correct.

b) 2(-2 + w) = 18Use the distributive property to expand 2(-2 + w). 2(-2) + 2(w) = 18-4 + 2w = 18Add 4 to each side. -4 + 2w + 4 = 18 + 42w = 22Divide each side by 2.  $\frac{2w}{2} = \frac{22}{2}$ w = 11To verify the solution, substitute w = 11 into 2(-2 + w) = 18. Left side = 2(-2 + w)Right side = 18= 2(-2 + 11)= 2(9)= 18 Since the left side equals the right side, w = 11 is correct.

\_\_\_\_\_ 1. What operations would you use to solve each equation? **a)** -5h + 4 = 6First \_\_\_\_\_, then \_\_\_\_\_. **b)** 2 + 5p = -3First \_\_\_\_\_, then \_\_\_\_\_. 2. Solve, then verify the equation. 2(t - 1) = 12Use the distributive property  $2(\_) - 2(\_) = 12$ to expand 2(t - 1). Substitute  $t = \_$  into the equation. Left side = 2(t - 1)Right side = = 2( ) = \_\_\_\_\_ = Since the left side equals the right side, t = is correct.

### Practice

**1.** Solve each equation.

<b>a)</b> z + 9 = 10	<b>b)</b> $s - 4 = -12$
<b>c)</b> 6 + c = 2	<b>d)</b> 5 = v - 2

**2.** For each statement, write then solve an equation to find the number. Verify the solution.

a)	A number divided by 4 is $-3$ .	<b>b)</b> Three times a number is 15.
	= -3	= 15
	Left side =	
		Right side =
	Right side = $n = \$ is correct.	x =  is correct.

- **3.** Emma tried to solve the equation 4x = 16 by subtracting 4 from each side. Show the correct way to solve the equation. 4x = 16
- **4.** Solve each equation. Verify the solution.

\_\_\_\_\_

a)	5k - 6 = 24	<b>b)</b> $3 + 4y = -9$
	Left side = $5k - 6$	Left side = $3 + 4y$
	= 5() - 6	=
	=	=
	Right side $k = \$ is correct.	Right side $=$ y = is correct.


#### **5. a)** Tuyen tried to solve the equation 3x - 6 = 15 like this:

 $\frac{3x}{3} - 6 = \frac{15}{3}$  x - 6 = 5 x - 6 + 6 = 5 + 6 x = 11Where did she make a mistake?

**b)** Show the correct way to solve 3x - 6 = 15. Verify the solution.

	Left side = $3x - 6$	Right side =	-
	= = Since the left side equals the		
<b>6.</b> A <i>w</i>	rectangle has length 4 cm an	d perimeter 12 cm.	The perimeter is the sum of all the sides.
	Write an equation that can = = Solve the equation.		dth of the rectangle.
c)	The width is cm. Verify the solution. Left side =	Right side =	-

# 6.2 Skill Builder



### Check

**1.** Use algebra tiles to solve: 4m + 6 = -2Record your steps algebraically.


# 6.2 Solving Equations by Using Balance Strategies

# **FOCUS** Model a problem with a linear equation, use balance strategies to solve the equation pictorially, and record the process symbolically.

To solve an equation, isolate the variable on one side of the equation.


We can use balance scales to model this.

Everything we do to one side of the equation must be done to the other side. This way, the scales remain balanced.




**b)** Check: Substitute x = 2 in each side of the equation. Left side = 3x + 2 = 3(2) + 2 = 6 + 2 = 8Since the left side equals the right side, x = 2 is correct.


# Check




If we have an equation with negative terms, it is easier to use algebra tiles to model and solve the equation.

We add the same tiles to each side or subtract the same tiles from each side to keep the equation balanced.





- 1. Use algebra tiles to model and solve the equation.
  - Record your work algebraically.



# **Example 3** Solving Equations with Rational Coefficients

Solve the equation, then verify the solution.

$$\frac{2a}{3} = 6$$

## Solution

Create an equivalent equation without fractions.

To clear the fraction, multiply each side by the denominator.


 $\frac{2a}{3} = 6$ Multiply each side by 3.  $\frac{2a}{3} \times 3 = 6 \times 3$   $\frac{2a}{3} \times \frac{3}{1} = \frac{2a}{1} = 2a$ Divide each side by 2.  $\frac{2a}{2} = \frac{18}{2}$  a = 9 Check: Substitute a = 9 in  $\frac{2a}{3} = 6$ . Left side  $= \frac{2a}{3}$  Right side = 6  $= \frac{2(9)}{3}$   $= \frac{18}{3}$  = 6Since the left side equals the right side, a = 9 is correct.


# Check

<b>1.</b> Sol <sup>y</sup>	ve. Verify the solution.	
a)	$\frac{\frac{x}{4}}{\frac{1}{2}} = 5$	Clear the fraction. Multiply each side by the denominator, 4.
	Check: Substitute <i>x</i> =	$_{-} in \frac{x}{4} = 5.$
	Left side =	Right side =
	=	
		x = is correct.
b)	$\frac{x}{4} + \frac{7}{4} = \frac{5}{4}$	
	Check: Substitute $x =$	$_{-}$ in $\frac{x}{4} + \frac{7}{4} = \frac{5}{4}$ .
	Left side =	Right side =
	Since	x =  is correct.

### Practice

**1.** Write the equation represented by the picture. Solve, and record your steps algebraically.







#### Can you ...


- Model a problem with a linear equation, and solve the equation pictorially and symbolically?
- Model a problem with a linear equation, use balance strategies to solve the equation pictorially, and record the process symbolically?

6.1 1. For each equation, what is the first operation you would do to isolate the variable?

a) 3k = 9(c) 2x - 3 = 4b) m - 2 = 5(c) 2x - 3 = 4b) m - 2 = 5(c) 5 = 3y - 4

2. For each statement, write then solve an equation to find the number. Verify the solution.

a) Two times a number is 10.	<b>b)</b> Three less than a number is 15.
<b>3.</b> Solve each equation.	
<b>a)</b> $x + 7 = -2$	<b>b)</b> 4c = 20
<b>c)</b> $4 = y - 2$	<b>d)</b> $\frac{m}{6} = 3$
<b>4.</b> Solve each equation. Verify the solution.	
<b>a)</b> 3q - 1 = 17	<b>b)</b> $2(3 + p) = -4$



**6.** Solve each equation. Verify the solution.

a)	3a - 2 = a - 6	<b>b)</b> $4 + h = 1 - 2h$	
	3a - 2 + 2 = a - 6 + 2	4 + h - 4 = 1 - 2h - 4	
	Left side = $3a - 2$	Left side = $4 + h$	
	=	=	_
	Right side = $a - 6$	Right side = $1 - 2h$	-
	=		_
			_
	a = is correct.	h =  is correct.	
c)	$\frac{5a}{6} = 10$		
<b>C</b> )	6		
		_	
		_	
	Left side = $\frac{5a}{6}$		
	-		
	=	-	
		-	
	Right side =	-	
	$a = \_$ is correct.		

# 6.3 Introduction to Linear Inequalities

#### **FOCUS** Write and graph inequalities.

Less than	<	below, under
Less than or equal to	$\leq$	up to, at most, no more than, maximum
Greater than	>	over, more than
Greater than or equal to	$\geq$	at least, minimum

## **Example 1** Writing an Inequality to Describe a Situation

Define a variable and write an inequality to describe the situation.



**b)** You must be at least 16 years old to get a driver's licence.


#### Solution

a) Let s represent the speed.You can go up to 60 km/h, but not faster.So, s can equal 60 or be any number less than 60.

The inequality is  $s \leq 60$ .

The inequality is  $a \ge 16$ .


b) Let a represent the age to get a driver's licence.
"At least 16" means that you must be 16, or older.
You cannot be less than 16.
So, a can equal 16 or be greater than 16.




# Check


- **1.** Let *t* represent the temperature in degrees Celsius. Write an inequality to describe each situation:
  - a) For temperatures less than 0°C, make sure to wear warm clothing. t \_\_\_\_\_ 0
- **b)** The highest temperature we've had this week was 12°C. *t* \_\_\_\_\_ 12

# **Linear Inequalities** A linear inequality may be true for many values of the variable.






Graph each inequality on a number line.
 Write 3 numbers that are possible solutions for each inequality.



#### **1.** Is each inequality true or false?

If it is false, change the sign to write a true inequality.



6. Write an inequality whose solution is graphed on the number line.



# 6.4 Solving Linear Inequalities by Using Addition and Subtraction


FOCUS Use addition and subtraction to solve inequalities.

Consider the inequality -1 < 3.

-3 -2 -1 0 1 2 3 4 5

What happens to an inequality if we add the same number to each side?

-1 < 3 Add 2 to each side. Left side: -1 + 2 = 1Right side: 3 + 2 = 5The resulting inequality is still true: 1 < 5




What happens to an inequality if we subtract the same number from each side? -1 < 3 Subtract 2 from each side.

Left side: -1 - 2 = -3

Right side: 3 - 2 = 1

The resulting inequality is still true: -3 < 1




#### **Property of Inequalities**


When the same number is added to or subtracted from each side of an inequality, the resulting inequality is still true.

The strategy that we used to solve an equation can be used to solve an inequality. Isolate the variable to solve.

Equation	Inequality
r - 6 = -2	r - 6 < -2
r - 6 + 6 = -2 + 6	r - 6 + 6 < -2 + 6
r = 4	<i>r</i> < 4
There is only 1 solution: $r = 4$	Any number less than 4 is part of the solution.
	The solution includes 3, 2, and 1, for example.



1. Solve each inequality. Graph the solution on the number line.



## **Example 2** Solving an Inequality with Variables on Both Sides

- **a)** Solve the inequality 3d + 2 < 2d 2.
- **b)** Graph the solution on a number line.

### Solution

a) 3d + 2 < 2d - 2 3d + 2 - 2d < 2d - 2 - 2d d + 2 < -2 d + 2 - 2 < -2 - 2d < -4


**b)**  $\leftarrow$  **b**  $\rightarrow$  **b**

Subtract 2*d* from each side.

Subtract 2 from each side.

Place an open circle on -4 because -4 is not part of the solution.

**1.** Solve each inequality. Graph the solution on a number line.



\_\_\_\_\_

#### Practice

**1.** Which operation will you perform to each side of the inequality to isolate the variable?


<b>a)</b> a + 1 > 3	<b>b)</b> 2 < m - 3
<b>c)</b> $x - 4 \ge 5$	<b>d)</b> 6 > 1 − z


- **2.** Fill in the missing steps to get to the solution.
  - **a)** x + 5 > 10  $x + 5 \_ > 10 \_$   $x > \_$  **b)**  $12 \le x - 4$   $12 \_ \le x - 4$  $\_ \le x$

#### 3. Solve each inequality.


Match each inequality with the graph of its solution, below.

a)	n - 4 > -2	<b>b)</b> p	+ 6 < -2
c)	$u-3 \ge -4$	<b>d)</b> 2	+ y > -2
	i) <u>-2 -1 0 1 2</u>	<u>+</u> →	
	<b>ii)</b> + + + + + + + + + + + + + + + + + + +	<u> </u>	
	iii) <u>+ + + + + + + + + + + + + + + + + + +</u>	<u>↓</u> →	
	<b>iv)</b> ← + + + ← -12 -11 -10 -9 -8		






# 6.5 Solving Linear Inequalities by Using Multiplication and Division





What happens to an inequality when we multiply or divide each side by the same negative number?



#### **Property of Inequalities**

When each side of an inequality is multiplied or divided by the same negative number, the inequality sign must be reversed for the inequality to remain true.





**1.** State whether you would reverse the inequality sign to solve each inequality.

a)	-2 <i>m</i> < 8	<b>b)</b> 2 <i>m</i> ≤ 8	<b>c)</b> $\frac{y}{-2} > 3$
<b>7</b> Solv	ve the inequalities in questior	n 1. Graph each solution	
	-2m < 8	<b>b)</b> $2m \le 8$	<b>c)</b> $\frac{y}{-2} > 3$
,		<i>.,</i>	-2
	 _5 _4 _3 _2 _1 0		

Example 2 Solving a Multi-Step Inequality **a)** Solve the inequality:  $1 - \frac{2}{3}x > 3$ **b)** Graph the solution. Solution ----**a)**  $1 - \frac{2}{3}x > 3$ Subtract 1 from each side to isolate *x*.  $1 - \frac{2}{3}x - 1 > 3 - 1$  $-\frac{2}{3}x > 2$ Multiply each side by -3 to clear the fraction. Reverse the inequality sign.  $(-3)\left(-\frac{2}{3}x\right) < (-3)(2)$ 2x < -6Divide each side by 2.  $\frac{2x}{2} < \frac{-6}{2}$ x < -3

**b)** The solution of x < -3 is all numbers less than -3.

-7 -6 -5 -4 -3 -2

#### Check

**1.** Solve the inequality:  $-\frac{2f}{5} < 4$ 

Graph the solution on the number line.

 -+-+-+-+	If you multiply or divide by a negative number, remember to reverse the inequality sign.

Practice

- **1.** a) Will the inequality sign change when you perform the indicated operation on each side of the inequality?
  - i) 3 > -2; Multiply by 2
  - ii)  $4 \le 8$ ; Divide by -4
  - iii) -5 < 1; Multiply by -5
  - **iv)** 1 > -4; Divide by 1

**b)** Perform each operation above. Write the resulting inequality.


- i) ii) \_\_\_\_\_ iii) iv)
- **2. a)** For the inequality -2 < 6, identify which of the following operations will reverse the inequality sign.
  - i) Multiply both sides by -4ii) Divide both sides by 2
  - **b)** Perform each operation above. Write the resulting inequality. i) \_\_\_\_\_
    - ii) \_\_\_\_\_

#### 3. a) What operation do you have to do to solve each inequality?

- i) 3*x* > 9 **ii)** -4p < -8
  - **iii)**  $-3y \le 15$

**iv)**  $\frac{q}{-2} \le 5$ 

**b)** State whether you would reverse the inequality sign to solve each inequality in part a.



# Unit 6 Puzzle

#### How Great Is My Number? You will need

10 red tiles, 10 yellow tiles, 2 die Label 1 die with the following faces:

> < = 2 ≤	=
-----------	---

#### Number of Players

#### 2

#### Goal of the Game

To get 4 tiles in a row, vertically, horizontally, or diagonally

8	-1	0	2	-2
-4	9	-5	4	1
3	7	6	5	-9
-12	-10	-7	-3	11
-6	-8	10	12	-11

#### How to Play

**1.** Roll the number die.

The player with the greater number goes first.

- **2.** The starting player rolls the die, and covers a number on the board that corresponds to what was rolled. For example, if a player rolls  $\geq$   $\bullet$ , the player can cover any number that is greater than or equal to 1.
- **3.** Only 1 number can be covered on a turn.
- 4. Players alternate turns.
- 5. The first player to get 4 tiles in a row wins.

# Unit 6 Study Guide

Skill	Description	Example
Solving Equations	To solve an equation, find the value of the variable that makes the left side of the equation equal to the right side.	Solve the equation: 3y - 2 = y + 4 Solution
	To solve an equation, isolate the variable on one side of the equation.	3y - 2 = y + 4 3y - 2 + 2 = y + 4 + 2 3y = y + 6 3y - y = y - y + 6
	<ul> <li>Use inverse operations or a balance strategy to perform the same operation on both sides of the equation:</li> <li>Add the same quantity to each side</li> <li>Subtract the same quantity from each side</li> <li>Multiply or divide each side by the same non-zero quantity</li> </ul>	$2y = 6$ $\frac{2y}{2} = \frac{6}{2}$ $y = 3$
	Algebra tiles and balance scales can help model the steps in the solution.	
Solving Inequalities	An inequality is a statement that one quantity is less than (<) another, greater than (>) another, less than or equal to ( $\leq$ ) another, or greater than or equal to ( $\geq$ ) another.	Solve the inequality and graph the solution: $-2s - 2 \le s - 5$ Solution $-2s - 2 + 2 \le s - 5 + 2$
	The inequality sign reverses when you multiply or divide each side of the inequality by the same negative number.	$-2s \le s - 3$ $-2s - s \le s - 3 - s$ $-3s \le -3$ $\frac{-3s}{-3} \ge \frac{-3}{-3}$ $s \ge 1$
	A linear inequality may be true for many values of the variable. We can graph the solutions on a number line.	Since we divide each side by the same negative number, the inequality sign is reversed. -+ $+$ $+$ $+$ $+$ $+$ $+$ $+$ $+$ $+$

# **Unit 6 Review**

**6.1 1.** Solve each equation. Verify the results. **a)** f + 6 = 3**b)** g - 5 = -2 $f = \_$  is correct.  $g = \_$  is correct. **d)** -2k = 6**c)** 5*h* = 25  $h = \_$  is correct.  $k = \_$  is correct. 2. Solve each equation. Verify the solution. **b)** 2 - 3c = -7**a)** 4x - 2 = 6c =\_\_\_\_\_ is correct. x =\_\_\_\_\_ is correct. **c)** 2v - 3 = -9**d)** -2(2 + w) = -20v =\_\_\_\_\_ is correct. w =\_\_\_\_\_ is correct.

**6.2 3.** Write the equation modelled by each set of algebra tiles. Solve the equation.

a)	b)
<b>4.</b> Solve each equation.	
<b>a)</b> $9 - 2w = w - 6$	<b>b)</b> $e - 6 = 6 - e$
<b>5.</b> Solve each equation. Verify the solution <b>a)</b> $6 + \frac{s}{2} = 7$	on. Left side = $6 + \frac{s}{2}$
	Right side = $s = \$ is correct.

<b>b)</b> $4 + \frac{2x}{3} = 2$	Left side = $4 + \frac{2x}{3}$
	Right side =
	x =  is correct.
<b>6.3 6.</b> Graph each inequality. Write 3 numbers that are possible solution	ons for each inequality.
<b>a)</b> $q > -3$ $+$ $+$ $+$ $+$ $+$ $+$ $+$ $+$ $+$ $+$	<b>b)</b> $w \le 0$ -+-+++++++++++++++++++++++++++++++++
<b>c)</b> $t \ge -1$ + + + + + + + + + + + + + + + + + + +	<b>d)</b> <i>r</i> < 6 ++++++++++++++++++++++++++++++++++
<b>7.</b> Write an inequality whose solution is gra	aphed on the number line.
<b>a)</b> -3 -2 -1 0 1	<b>b)</b> 1 2 3 4 5
<b>6.4 8.</b> Solve each inequality. Graph the solution	۱.
<b>a)</b> d-6>4	<b>b)</b> 2f + 1 < -3
9 10 11 12 13	
9. Solve each inequality. Graph the solution	۱.
<b>a)</b> 4 <i>j</i> − 1 ≥ 2 <i>j</i> + 3	<b>b)</b> <i>k</i> − 2 < 2 − <i>k</i>

6.5 10.State whether you would reverse the inequality sign to solve each inequality.

